

GARDASIL is the only vaccine that may help guard against diseases that are caused by HPV Types 6, 11, 16, and 18:

- Cervical cancer
- Cervical abnormalities that can sometimes lead to cervical cancer
- Genital warts

HPV Types 16 and 18 cause 70% of cervical cancer cases, and HPV Types 6 and 11 cause 90% of genital warts cases.

IMPORTANT INFORMATION ABOUT GARDASIL

GARDASIL will not treat these diseases and will not protect against diseases caused by other types of HPV. GARDASIL may not fully protect everyone.

Anyone who is allergic to the ingredients of GARDASIL should not receive the vaccine.

GARDASIL can cause pain, swelling, itching, and redness at the injection site, fever, nausea, and dizziness. GARDASIL is not for women who are pregnant.

PROTECTION: 3 IS KEY

GARDASIL is given as 3 injections over 6 months.

- First dose: at a date you and your daughter's doctor or healthcare professional choose
- Second dose: 2 months after the first dose
- Third dose: 6 months after the first dose

If your daughter's doctor or healthcare professional decides GARDASIL is right for her, make sure she gets all 3 doses. This will allow her to get the full benefits of GARDASIL.

GARDASIL AND YOUR DAUGHTER

GARDASIL works when given before your daughter has any contact with HPV Types 6, 11, 16, and 18. That's why getting your daughter vaccinated with GARDASIL today may help guard her against cervical cancer and other diseases caused by HPV in the future.

IMPORTANT INFORMATION ABOUT GARDASIL

GARDASIL is not for women who are pregnant. Anyone who is allergic to the ingredients of GARDASIL should not receive the vaccine. GARDASIL may not fully protect everyone, and GARDASIL does not prevent all types of cervical cancer, so future cervical cancer screenings will be important for your daughter. GARDASIL can cause pain, swelling, itching, and redness at the injection site, fever, nausea, and dizziness.


GARDASIL.
[Quadrivalent Human Papillomavirus
(Types 6, 11, 16, 18) Recombinant Vaccine]